


Relieves hot flashes from menopause and strengthens your bones. Duavee (conjugated estrogens / bazedoxifene) is a once daily pill that helps with symptoms of menopause.

What to expect when starting

Here's when people typically experience benefits as well as some possible side effects. Your experience may be different.



† Data on frequency of side effects is not available.

Tips from pharmacists

- Duavee (conjugated estrogens / bazedoxifene) comes in a blister pack that is sealed in a foil pouch. Don't push the pills out of the pack until you are going to take your dose. Don't use a pill organizer to store this because it is important to protect the tablet from moisture.
- Write the date you open the foil pouch on the blister pack. You cannot use the pills if more than 60 days have passed since you opened the foil pouch.
- Take 1 pill at the same time every day. Don't double up on pills if you miss a dose.
- You can take Duavee (conjugated estrogens / bazedoxifene) at the same time as calcium and vitamin D supplements if you are taking them.
- If you need to have surgery, you must stop taking Duavee (conjugated estrogens / bazedoxifene) at least 4 to 6 weeks before the surgery is performed.
- Tell your doctor right away if you notice unusual vaginal bleeding.

Risks and warnings

The FDA has identified these as possible serious or life-threatening risks with this medication. Contact your doctor if you experience any of these.

Endometrial cancer

Using medicines like Duavee (conjugated estrogens / bazedoxifene) for more than a year can increase your risk of uterus (endometrial) cancer. Let your doctor know if you notice any unusual bleeding. Talk to your doctor about other medicines you can take with Duavee (conjugated estrogens / bazedoxifene) to protect yourself from cancer if you are worried.

Increased risk of blood clots and stroke

Duavee (conjugated estrogens / bazedoxifene) can increase your risk of blood clots, which can be very dangerous. You're more likely to have them if you smoke tobacco, are obese, or if you or your family have ever had blood clots. Get medical care right away if you have pain in your legs or arms, bad headaches, trouble breathing, or sudden changes in your speech or vision.

Increased risk of heart disease

Medicines with estrogen like Duavee (conjugated estrogens / bazedoxifene) can increase your risk of heart disease, and this risk gets worse the longer you take it. Talk to your doctor regularly to see if you still need to use Duavee (conjugated estrogens / bazedoxifene).