

Edarbyclor (azilsartan / chlorthalidone) basics

Lowers blood pressure. Edarbyclor (azilsartan / chlorthalidone) is a good 2-in-1 blood pressure-lowering medicine, especially if you are at high risk for stroke and heart attacks.

What to expect when starting

Here's when people typically experience benefits as well as some possible side effects. Your experience may be different.

	Hours	Days	Weeks	Months	Long term
Edarbyclor (azilsartan / chlorthalidone) Drug effectiveness					
Possible side effects					
Dizziness 9% chance					
Fatigue 2% chance					
Low blood pressure 2% chance					
Fainting†					

† Data on frequency of side effects is not available.

Tips from pharmacists

- Take with or without food.
- Avoid driving and doing other tasks that require you to be alert until you know how this drug affects you.
- Get your blood tests regularly and as told by your doctor.
- Edarbyclor (azilsartan / chlorthalidone) increases your urination. Take your dose in the morning or a few hours before bedtime if you find yourself waking up in the middle of the night to go to the bathroom.

Risks and warnings

The FDA has identified these as possible serious or life-threatening risks with this medication. Contact your doctor if you experience any of these.

Harm to fetus

Edarbyclor (azilsartan / chlorthalidone) causes birth defects. If you're pregnant, stop taking Edarbyclor (azilsartan / chlorthalidone) right away and talk to your doctor.

Low blood pressure

Edarbyclor (azilsartan / chlorthalidone) lowers your blood pressure. Changes in blood pressure can make you feel lightheaded or dizzy. Stand up slowly, especially when you have been sitting or laying down for a while.

Worsening kidneys

Edarbyclor (azilsartan / chlorthalidone) can impair your kidney function, especially if you have pre-existing kidney disease. Your doctor will order blood tests regularly to monitor your kidneys.