


Odefsey (emtricitabine / rilpivirine / tenofovir) basics

Powered by
IODINE

Treats HIV infection Odefsey (emtricitabine / rilpivirine / tenofovir) is a convenient 3-in-1 medication used to treat HIV infections. It's similar to Complera, but contains a newer HIV drug which has shown to cause less harmful effects on your kidneys and bones.

What to expect when starting

Here's when people typically experience benefits as well as some possible side effects. Your experience may be different.



† Data on frequency of side effects is not available.

Tips from pharmacists

- Odefsey (emtricitabine / rilpivirine / tenofovir) should be taken at the same time everyday.
- Take each dose with a light meal.
- Talk to your doctor immediately if you notice yellowing of eyes or skin, dark urine, rash, fever, fatigue, or have difficulty breathing, as these may be signs of liver problems or sensitivity reactions.

Risks and warnings

The FDA has identified these as possible serious or life-threatening risks with this medication. Contact your doctor if you experience any of these.

Worsening hepatitis B infection

If you have hepatitis B while taking Odefsey (emtricitabine / rilpivirine / tenofovir), stopping Odefsey (emtricitabine / rilpivirine / tenofovir) can worsen the hepatitis infection. Your doctor will test you before and during HIV treatment to see if you have hepatitis B. If you do have it, your doctor will provide regular blood tests to make sure your liver is ok, and might put you on medicines for your hepatitis.

Lactic acidosis | Enlarged, fatty liver

Odefsey (emtricitabine / rilpivirine / tenofovir) can cause this rare but life-threatening condition. Symptoms include difficulty breathing, nausea, vomiting, right sided stomach pain, and dark urine . This is a medical emergency and you should get help right away.

Mood and behavior changes

Odefsey (emtricitabine / rilpivirine / tenofovir) has been associated with depression, thoughts of suicide, negative thoughts, and depressed mood. If your mood starts to change while taking Odefsey (emtricitabine / rilpivirine / tenofovir), let your doctor know. If you start thinking about wanting to harm yourself, seek medical attention immediately.