


Sanctura (trospium) basics

Treats overactive bladder. Sanctura (trospium) is a good treatment for overactive bladder after exercises when other methods to control it haven't worked.

What to expect when starting

Here's when people typically experience benefits as well as some possible side effects. Your experience may be different.



* Placebo is a sugar pill with no medicinal benefit.

† Data on frequency of side effects is not available.

Tips from pharmacists

- Take Sanctura (trospium) on an empty stomach 1 hour before your meal with a glass of water.
- You are more likely to have heat stroke on Sanctura (trospium). Drink plenty of water, stay out of the sun, and don't exercise too hard if it's hot out.
- Can cause dry mouth, constipation or headache.
- Sanctura (trospium) can make you dizzy or cause vision changes, so don't drive until you know how it affects you.
- If you miss a dose, wait until 1 hour before your next meal to take your next dose. Don't take extra medicine to make up for the missed dose.

Risks and warnings

Contact your doctor if you experience any of these.

Facial swelling & breathing problems

Sanctura (trospium) may cause an uncommon, but serious side effect called angioedema, or swelling of the face, lips, tongue, or throat. This can make it difficult to breathe. Go to the emergency room right away if you suddenly can't breathe or notice swelling around your face or throat.

Difficulty concentrating

Sanctura (trospium) can make you less alert during the first few days you take it. Avoid doing things that require you to focus, such as driving a car. Drinking alcohol or taking medicines that also make you lose focus with Sanctura (trospium) can make this worse.

Vision changes

Sanctura (trospium) may cause vision changes such as blurred vision or worsening of glaucoma symptoms. If you have glaucoma, talk to your doctor before using Sanctura (trospium).